

Mathis Bloßfeld, Horst Müller, Detlef Angermann

Deutsches Geodätisches Forschungsinstitut, Munich, Germany, blossfeld@dgfi.badw.de

Motivation

Satellite Laser Ranging (SLR) is the primary technique to estimate consistently station positions, Earth Orientation Parameters (EOP) and orbit parameters of the satellites together with the spherical harmonics of low degree and order of the Earth gravity field. The big effort of the common adjustment of these parameters is the high correlation of the orbit parameters (e.g. Kepler elements, empirical accelerations), length of day (LOD) as the first derivative of Universal Time (UT) and the gravity field parameter C_{20} . The relation between these parameters is given in equation (1).

$$\dot{\Omega} \Big|_{\text{secular}} = \frac{3}{2} \sqrt{\frac{GM}{a_e^3}} \left(\frac{a_e}{a}\right)^{\frac{7}{2}} \frac{C_{20} \cos i}{(1-e^2)^2}, \quad (1)$$

where $\dot{\Omega}$ is the rate of change of the ascending node, a_e is the major axis of the Earth, GM is the gravity constant multiplied by the mass of the Earth and a, e, i are the major axis, the eccentricity and the inclination of a satellite.

In this study we discuss different solutions (7-day and 28-day arc, one-satellite and multi-satellite constellation) and evaluate the correlations and the stability of the estimated parameters.

Solution types




Fig. 1: Fits of the 7-day/28-day orbits of LAGEOS 1 and 2. Only observations from official core stations of the International Laser Ranging Service (ILRS) are considered.

Fig. 1 shows the different fits of the satellite orbits of LAGEOS 1 and 2. The mean RMS values are given in Tab. 1. The accuracy of the 7-day arcs is at the level of five millimeters whereas the accuracy of the 28-day arcs is at the level of one centimeter. Fig. 2 points out that a reduction of the correlation of C_{20} and Ω could be achieved using longer arcs or including more than one satellite. The most uncorrelated parameters could be estimated in solutions containing two satellites with an arc length of 28 days.




Fig. 2: Mean correlations of C_{20} and Ω for one-satellite solutions (green-coloured) and multi-satellite solutions (orange-coloured). In the left part of Fig. 2 the mean correlations of solutions with an arc length of 7 days are shown, whereas the right part illustrates the same situation for a 28-day arc.

Earth Orientation Parameters (1)

The parameterization of UT1 and LOD is the same in all solutions. Since SLR is not able to determine UT1, the offsets are extrapolated with LOD to 0h epochs of a piecewise linear polygon. At the mid-epoch of the arc, one UT1 value is fixed to a priori (IERS 08 C04).

Because of the high correlations expressed in equation (1), errors or non-modelled perturbations of the satellites systematically affect the estimated LOD and the UT1 values respectively (Fig. 3).




Fig. 3: Accumulated differences of UT1 w.r.t. IERS 08 C04 over a time span of 16.5 years. The individual solutions of LAGEOS 1 and 2 with an arc length of 28 days are not displayed (see Tab. 2).

Fig. 3 shows a systematic drift for UT1 w.r.t. the IERS 08 C04 time series. Tab. 2 summarizes the different mean drifts for each solution.

	LAGEOS 1	LAGEOS 2	LAGEOS 1&2
7-day arc	8.23	-17.57	-3.63
28-day arc	-38.02	-26.93	-3.97

Tab. 2: Mean drifts of UT1 [ms/y] w.r.t. IERS 08 C04 for the different solutions.

The spurious drifts of the 7-day arc one-satellite solution have an opposite sign and a nearly constant ratio which could be explained with equation (1). Since all parameters except the inclination of LAGEOS 1 and 2 are approximately the same, the sign and ratio depends on the $\cos i$ term of equation (1). The one-satellite solution with a 28-day arc doesn't show this characteristics. Although the mean correlation of C_{20} and Ω is reduced by using a 28-day arc multi-satellite solution (Fig. 2), there still remains a systematic drift in Fig. 3.

Gravity field parameter

The estimated gravity field coefficients of the solutions with the data of two satellites show a very good agreement with a solution of the Center for Space Research (CSR) although the CSR solution contains observations to three more satellites than the DGFI solutions (Fig. 4).




Fig. 4: Estimated normalized C_{20} coeffs. The DGFI solutions contain only data from LAGEOS 1 and 2 whereas the solution of CSR includes in addition data from Stella, Starlette and Ajisai.

Earth Orientation Parameters (2)

The estimated gravity field coefficients of the multi-satellite solutions (Fig. 4) are then, in a second iteration step, set up as a priori values for C_{20} to reduce the drift of the UT1 values resulting from a slightly wrong C_{20} . The results are summarized in Fig. 5 and Tab. 3. The one-satellite solutions benefit tremendously in this second iteration step. The mean correlation of C_{20} and Ω is reduced to 0.05 for all solution types.




Fig. 5: Accumulated differences of UT1 w.r.t. IERS 08 C04 over a time span of 16.5 years. All solutions of the second iteration show a systematic drift in the order of -2.8 to -3.9 ms/y (Tab. 3).

Nevertheless, Fig. 5 and Tab. 3 show that all solutions contain a spurious drift w.r.t. IERS 08 C04. This could be due to the fact that an offset of LOD could also be caused by a periodically occurring perturbation perpendicular to the orbit plane of the satellite (cross track direction).

	LAGEOS 1	LAGEOS 2	LAGEOS 1&2
7-day arc	-2.78	-3.23	-3.62
28-day arc	-3.55	-3.87	-3.70

Tab. 3: Mean drifts of UT1 [ms/y] w.r.t. IERS 08 C04 for the different solutions.

Conclusions & Future Work

The main part of the UT1 drifts in Fig. 3 are induced by the fact that the a priori C_{20} values of the first iteration step (here GGM02S, see Fig. 4) leads to a wrong $\dot{\Omega}$ and as a consequence of this to a wrong LOD. If we use the C_{20} values of a multi-satellite solution (see also Fig. 4) instead of that we get much lower drifts (Fig. 5). These remaining drifts have the same sign and therefor couldn't be excited by a wrong C_{20} . The next step would be to study the relationship between perturbations offending the satellite in cross track direction. To improve the solution furthermore we want to introduce other geodetic satellites like Etalon 1 and 2, Stella, Starlette and Ajisai. We also want to estimate variance factors in the combination of different satellites in order to improve the relative weighting.

References:

- Beutler G., "Methods of Celestial Mechanics Volume 1&2", Springer, Berlin, 2005.
- Tapley et al., "GGM02 – An improved Earth gravity model from GRACE", Journal of Geodesy, Vol. 79, No. 8, pp 467-478, November 2005, DOI: 10.1007/S00190-005-0480-Z
- Pearlman, M.R., Degnan, J.J., and Bosworth, J.M., "The International Laser Ranging Service", Advances in Space Research, Vol. 30, No. 2, pp. 135-143, July 2002, DOI:10.1016/S0273-1177(02)00277-6.

Acknowledgement:

The authors want to thank the German Research Foundation DFG for funding the project within the research group FOR 584 "Earth Rotation and Global Dynamic Processes".